

Quantum theory as a statistical theory under symmetry.

Inge S. Helland

Department of Mathematics, University of Oslo
POBox 1053 Blindern, N-0316 Oslo, Norway.
E-mail: ingeh@math.uio.no

Abstract

Both statistics and quantum theory deal with prediction using probability. We will show that there can be established a connection between these two areas. This will at the same time suggest a new, less formalistic way of looking upon basic quantum theory.

A total parameter space Φ , equipped with a group G of transformations, gives the mental image of some quantum system, in such a way that only certain components, functions of the total parameter ϕ can be estimated. Choose an experiment/ question a , and get from this a parameter space Λ^a , perhaps after some model reduction compatible with the group structure. As in statistics, it is important always to distinguish between observations and parameters, in particular between (minimal) observations t^a and state variables (parameters) λ^a . Let $\mathbf{K}^a$ be the L^2 -space of functions of t^a , and let $\mathbf{H}^a$ be the image of $\mathbf{K}^a$ under the expectation operator of the model. The measure determining the L^2 -spaces is the invariant measure under the maximal subgroup which induces a transformation on Λ^a .

There is a unitary connection between $\mathbf{K}^a$ and $\mathbf{H}^a$, and then under natural conditions between $\mathbf{H}^a$ and $\mathbf{H}^b$ for different a and b . Thus there exists a common Hilbert space $\mathbf{H}$ such that $\mathbf{H}^a$ equals $U^a\mathbf{H}$ for a unitary U^a . In agreement with the common formulation of quantum mechanics, the vectors of $\mathbf{H}$ are taken to represent the states of the system. The state interpretation is then: An ideal experiment $\mathbf{E}^a$ has been performed, and the result of this experiment, disregarding measurement errors, is that the parameter λ^a is equal to some fixed number λ_k^a . This essentially statistical construction leads under natural assumptions to the basic axioms of quantum mechanics, and thus implies a new statistical interpretation of this traditionally very formal theory. The probabilities are introduced via Born's formula, and this formula is proved from general, reasonable assumptions, essentially symmetry assumptions.

The theory is illustrated by a simple macroscopic example, and by the example of a spin $\frac{1}{2}$ particle. As a last example we show a connection to inference between related macroscopic experiments under symmetry.

1 Introduction

Nancy Cartwright [4] has argued that physical laws are about our models of reality, not about reality itself. At the same time she argues (p. 186) that the interpretation of quantum mechanics should be seen entirely in terms of transition probabilities. The present paper is in agreement with both these statements. I will supplement the first statement, however, by saying a few words about models. For many reasonably complex phenomena, several models, and several ways to give a language for model formulation, can be imagined. In some instances these models, while appearing different, are so closely related that they give the same predictions about reality. If this is the case, I think many would agree with me that we should choose the model which has the most intuitive interpretation, also in cases where there may be given strong historical and culturally related arguments for other models. It may of course be the case that the conventional model is more suitable for calculations, but this should not preclude us from using more intuitive points of view when arguing about the model and when trying to understand more complex phenomena.

In the present paper I try to offer such a completely different, in my view more intuitive, modelling approach to quantum mechanics, and I show that the ordinary quantum formalism, at least the time-independent aspects of it, follows from it.

Another purpose of this paper is to argue for a logical connection between quantum theory and a natural extension of statistical theory, an extension where symmetry aspects are taken into account by defining group transformations on the state space (or parameter space in statistical terminology), and where complementarity may be introduced by defining a total parameter space to which several potential experiments may be coupled. These different experiments may each be assumed to be limited by a context which is such that only a part of the total parameter may be estimated.

My aim is to arrive at ordinary quantum mechanics from this extended statistical theory. From this one may formulate the tentative conclusion that at least some of the differences between the two sciences, the way the theories are formulated today, may have some cultural origin.

To start such a programme, we have to relate it to conventional quantum theory. The basic formalism of quantum theory is given in slightly different ways in different books, although all agree about the foundation. For definiteness I will consider the following three axioms, taken from Isham [12]. In the last Section of the paper I will give a derivation of them (and in fact more) under suitable assumptions from the setting indicated above.

Rule 1.

The predictions of results of measurements in quantum mechanics made on an otherwise isolated system are probabilistic in nature. In situations where the maximum amount of information is available, this probabilistic information is represented mathematically by a vector in a complex Hilbert space $\mathbf{H}$ that forms the *state space* of the quantum theory. In so far as it gives the most precise

predictions that are possible, this vector is to be thought of as the mathematical representation of the physical notion of ‘state’ of the system.

Rule 2.

The observables of the system are represented mathematically by self-adjoint operators that act on the Hilbert space $\mathbf{H}$.

Rule 3.

If an observable quantity $\hat{\lambda}^a$ is represented by the self-adjoint operator T^a , and the state by the normalized vector $v \in \mathbf{H}$, then the expected result of the measurement is

$$E_v(\hat{\lambda}^a) = v^\dagger T^a v. \quad (1)$$

These three rules together with the Schrödinger equation constitute according to [12] the basic axioms of quantum mechanics. In this paper I will indicate the derivation of these rules from a setting which in essence is a natural extension of statistical theory. I will also comment upon issues like superselection rules and the extension to mixed states, and try to argue that formal equations like (1) can be associated with a very natural interpretation. A basic step in the derivation of the three rules is to give arguments for (a variant of) Born’s celebrated formula for transition probabilities in quantum mechanics.

Note that the states of quantum mechanics, in the non-degenerate case, can be interpreted in the following way: Corresponding to the operator T there is a physical variable λ , and a state, as given by a state vector v_k is connected to a particular value λ_k of λ . Thus we can say that the state is determined by two elements: 1) A question: What is the value of λ ? 2) The answer: $\lambda = \lambda_k$. In this paper we show that we can go the other way, and start with a question/answer pair like above, and then under certain assumptions, mainly related to symmetry and to a limited context, we arrive at the Hilbert space formulation with the above interpretation of the state vectors.

Consider the spin 1/2 particle, and let a be any 3-vector. Then it makes sense to say: Question: What is the spin component in direction a ? Answer: +1. And to say that these two elements together define a state, in ordinary quantummechanical language as the state with eigenvalue +1 for the operator $a \cdot \sigma$, where σ is the vector composed of the three Pauli spin matrices.

My aim is to approach this in a less formal way than the Hilbert space approach, the approach that we now have become so familiar with, but which seems to be impossible to explain in a simple way to people that have not obtained this familiarity. To carry out such a programme, I want to use other, more direct, mental models.

The point is that I consider the spin vector in the model as what I call a total parameter (this name was suggested to me by Peter Jupp). A total parameter is to me something that can be formulated by ordinary language and is associated by a tentative model of the subatomic reality, but for which it is

far too optimistic to connect a definite value in the sense that this value can be estimated by an experiment.

I will give examples of this below, and I will give more examples in the book [9].

What we can confront with experiments, though, is some given function of this spin vector, like a component of the spin vector in a certain direction, or more precisely, the sign of this component, say, in the spin $1/2$ case. The rest of the spin vector will always remain unknown to us. One might of course say that then it is nonsense to speak about the rest of the spin vector, but I would say that it is useful to have a mental model. Various people may have variants of the same mental model, but this does not matter as long as they agree about the symmetry aspect and about the observable part of the spin vector.

A relatively concrete realisation of this is given by the 'triangle in a sphere' which I describe below for which we are only allowed to look through one chosen window. Here we may have constructed the triangle with a given colour, made it from a definite material and so on, but to an observer in a window, all these parameters may only be imagined mentally. All that matters for his observations are the corners A, B or C , and all that matters in order to interpret these observations is a mental model of a triangle with rotational symmetry attached to it.

Or consider the case of a single patient at a fixed time, where we might be interested in expected recovery times τ_1 and τ_2 under two potential treatments (and in other parameters), so that the vector $\tau = (\tau_1, \tau_2)$ does not have an empirical value, but one component can be estimated by an experiment.

But I repeat: The total parameter may nevertheless be a useful quantity. In the model context it may help us to just have a mental picture of what we think is going on, say, in the subatomic world. The spin vector can be red or blue, can be imagined to be connected to some solid body, or just be an arrow. But what is the same in all these mental pictures, is every single component that we are allowed to ask questions about.

One very useful property of the total parameter is that we can imagine group transformations of it, and that these transformations then have consequences for the observable components. In the spin case and in the case with the triangle in the sphere, we can think of rotations. In the treatments of a patient-example it is meaningful to study scale invariance: $(\tau_1, \tau_2) \rightarrow (b\tau_1, b\tau_2)$.

So for such a total parameter I cannot ask every question I want, but I ask a question about a maximally observable component, and the answer is what I say defines a state. Even though I start with a mental picture involving unobservables, the state is defined in terms of an observable quantity.

Return to the spin $1/2$ case; there one can get an even more direct characterization: Let the question be about the spin in direction a , and let the answer be $+1$, then define as an abbreviation for this state, the 3-vector $u = a$. If the answer is -1 , let the state be characterized by the 3-vector $u = -a$. This is consistent, since the latter state also can be result $+1$. Thus from my definition, the state can be characterized by a vector u .

It is well known that in this case the state vectors v in the Hilbert space stand

(again disregarding an irrelevant phase factor) in one-to-one correspondence with the Bloch sphere operators $1/2(I + u \cdot \sigma)$, which again are in one-to-one correspondence with the vectors u , vectors which can be defined as in the previous paragraph. Thus the state can be characterized either by the Hilbert space vector v , the 3-vector u or the Bloch sphere operator.

Later I generalize this to mixed states and further to the effects of Busch [3] and Caves et al. [5], used below to prove the Born formula.

In the general Hilbert space every normalized vector is the eigenvector of some operator T , and if this corresponds to a meaningful physical variable, then you have a question and answer situation. In my approach there seem to be some uniqueness here under certain assumptions.

One of my hopes has been to find a meaningful link between quantum theory and statistical theory, and my feeling is that we are close to that here. But I also feel that we then must be willing to change focus slightly in traditional theoretical statistics.

In theoretical statistics there seems to be almost complete separation between experimental design theory and inference theory. In practical experiments the two are linked closely together. In fact, a formulation of the experimental question should almost always be an important part of the conclusion, and in any useful investigation in biology and medicine, say, it always is. Thus here also, in my opinion the conclusion should, in all good experiments, be stated as a question plus an answer.

People that are close to applications, sometimes think in terms of mental quantities that are close to my total parameters, for instance Searle in his book [14] bases his treatment to a large extent on unestimable linear parameters, motivated by the fact that this gives a nicer mental picture when you want to consider several models at the same time.

A more detailed derivation of the rules 1-3, together with more comments on several issues of quantum mechanics, also the Schrödinger equation, will be given in the forthcoming book [9]. There is also a paper [10] under preparation, intended for the statistical community.

2 The statistical background.

Statistics is a tool in almost any science which collects and analyses empirical data. In statistical theory it is very important to distinguish between state variables on the one hand, and observations on the other hand. In statistics the state variables are usually called *parameters*. This distinction is unfortunately not common to draw in theoretical physics, but in quantum mechanical papers there is much discussion related to measurement apparatus. When I now will formulate the standard statistical model, it can be useful to have such an apparatus in mind. I will return to this issue later.

First, it is useful in our setting to stress that every experiment has as its purpose to answer one or several questions about nature. Call such an experiment $\mathbf{E}^a$, and let a denote the set of questions that is posed in this particular

experiment. The ideal answer to these questions are the values of our state variables θ^a , or what statisticians call the parameter of experiment $\mathbf{E}^a$. This can for instance be the expected recovery time of patients with a particular disease from some given population that receive a certain treatment. The observations may then be the realized recovery times $y = (y_1, \dots, y_n)$ for n randomly selected patients from this population. In statistical inference theory and practice such observations act as measurements, if you like, by some measurement apparatus, where the purpose is to make statements about the unknown reality represented by θ^a , in particular, to find an estimate $\hat{\theta}^a$ from data.

This conceptual framework has turned out to be useful in a large number of sciences. It is the purpose of this paper to argue that it can be of some use in the process of finding a non-formal foundation of quantum physics, too. The measurement apparatus in physics can be perfect in the sense that they give the exact correct value, but nevertheless it is of value to distinguish between the observations, say positions of pointers or more generally some points in an Euclidean measure space, and the ideal values, here the parameters θ .

For a general non-ideal measurement apparatus, our statistical model says that, given the state of nature θ there is a probability distribution $Q^\theta(dy)$ for the observations y . The precise form of this distribution will not be important for us here, but in macroscopic experiments it is very important, and forms the basis for statistical inference: How to use uncertain observations to make statements concerning the state of nature, for example about expected recovery times of patients under some treatment.

One special concept from this setting is that of a *sufficient observation*: An observational function $t(y)$ is called sufficient if the conditional distribution under the model of y given t is independent of the parameter θ . This means that all information about the unknown parameter is contained in the reduced observation t , which may be very useful in some settings. In the simplest cases, for instance, of the recovery time example above, the mean recovery time $\bar{y} = \sum_i y_i/n$ is sufficient for the expected recovery time θ .

A sufficient (reduced) observation t is called *complete* whenever the following holds: If $h(t)$ has the property that the expectation under the model is 0 for all θ , then $h(t)$ is identically 0 (almost surely under the model). Often, when a sufficient observation has been reduced as much as possible, it will be complete.

These technical concepts are mentioned here because they are needed at a specific point of our argument below. Much more information about these concepts and related concepts can be found in advanced books in mathematical statistics like [13]. There are many intermediate book in mathematical statistics; one of them is [6].

We end the Section by discussing two extensions of the standard statistical model which are important in practice, and will be important in our approach to quantum theory, though they are in my mind insufficiently treated in the statistical literature.

First, we discuss the genuine choice of experiments which was mentioned in the beginning of this Section. In ordinary statistics this is related to the issue of experimental design, which has its own large literature. In quantum physics we

can think of the choice between measuring the position or the momentum of a particle, or of measuring the spin of an electron in one of all possible directions. In each case we have the choice between several possible experiments $\mathbf{E}^a$, and given each choice, we have a measurement apparatus, which in our terminology may be modelled by a standard statistical model $Q^{\theta^a}(dy)$. Here θ^a is the aspect of the state of nature which is related to our chosen experiment, say the ‘real’ position of the particle in question.

It will be useful for us to introduce a quantity ϕ which is called the *total parameter* of the system in question, and let the parameters of the single experiment be functions of this total parameter: $\theta^a = \theta^a(\phi)$. In a quantum mechanical setting, the total parameter ϕ will not take a value, in the sense that it cannot be estimated from any experiment, and this may in fact happen in a macroscopic situation also: Consider one particular patient at one point of time, and let θ^1 be his expected recovery time if he is given a treatment 1, and let θ^2 be his expected recovery time if he is given a treatment 2. The parameters θ^1 and θ^2 are both unknown here, but only one of these can be confronted with an experimental value. Thus $\phi = (\theta^1, \theta^2)$ can not be considered to have a value, except in a hypothetical sense. The reason why it nevertheless is useful to consider such a total parameter, will become clearer in the next Section.

Next: All the parameters that we have talked about have been conceptual. In some cases the description given by the parameter may be too detailed, again it may be impossible to give it a value, at least it is impossible to estimate it from any possible observation. One example of this was the total parameter ϕ above containing the result of mutually exclusive treatments. Another example can be $\theta = (\mu, \sigma)$ where μ is the ideal measurement value given by some fragile apparatus which is destroyed after one measurement, and where σ is some ideal measure of uncertainty, found by dismantling the apparatus, a process which also destroys it.

In such cases it may sometimes be helpful to introduce a reduced parameter $\lambda = \lambda(\theta)$, and it may then be the case that λ actually can be estimated through observations. In this paper we will then talk about a reduced model, a statistical model for the observations which only depends upon the parameter λ , not upon θ . In statistical practice such reduced models are sometimes chosen in order to improve prediction performance, even in cases where θ is estimable. In our setting the model reductions are necessary: Information beyond λ can never be found. Interestingly enough, there are statistical situations (collinearity in multiple regression) where connections between the two kinds of model reductions can be seen.

A somewhat different example of a parameter which do not take a value can be the expected drying time of the paint I use when painting my house, if the case is that I after some consideration choose not to paint my house at all. Such cases are related to the field of causal inference, where the concept of counterfactuals plays an important rôle.

As a final statistical concept, a *perfect* experiment $\mathbf{E}$ is an experiment where all measurement error may be disregarded, so that for the parameter λ , the estimate may be taken to be equal to the real value of the parameter: $\hat{\lambda} = \lambda$.

3 Symmetry in statistical models.

There is an important part of statistical theory which in addition to the statistical model assumes that there is a symmetry group acting upon the parameter space. The simplest such group may be induced by a change of units in the observations. More generally, for a scalar parameter we may have a scale group given by $\theta \mapsto b\theta$; $b > 0$, also written $\theta g = b\theta$, a location group given by $\theta \mapsto \theta + a$, written $\theta g = \theta + a$, or a scale and location group given by $\theta \mapsto b\theta + a$. Another important case is the rotation group for a vector parameter. For convenience, group actions will be written to the right of the parameter or observation to transform in this paper.

In general, a group acting upon the observations induces a group acting upon the parameters through using the statistical model [9, 10].

It is essential that the existence of such a transformation group G do not imply strong symmetry assumptions neither on the parameter space nor on the sample space. The main requirements are that the spaces should be closed under the group and that inference from $(yg, \theta g)$ should be as natural as inference from (y, θ) . This and other aspects are further discussed in [9].

For our purpose, it is important that a transformation group also can be defined upon a total parameter, even if this does not take a value. For example, in the case $\phi = (\theta^1, \theta^2)$, where θ^i is the expected recovery time for some fixed patient receiving treatment i , the time scale transformation given by $\phi g = (b\theta^1, b\theta^2)$ is definitively meaningful.

In this paper we will assume that the total parameter space Φ is locally compact and otherwise satisfies weak conditions which ensures that there is a right invariant measure ν under the group G on this space. This can be uniquely defined up to a multiplicative constant by first defining a right invariant measure ν_G on G by $\nu_G(Dg) = \nu_G(D)$; $D \subset G, g \in G$, and then taking $\nu(d(\phi_0 g)) = \nu_G(dg)$. In the corresponding case of an ordinary parameter, the right invariant measure can be recommended as a non-informative prior, a concept which is important in Bayesian statistics.

When Φ is a total parameter space with a transformation group G defined upon it, we may also be interested in the transformations on a parameter $\theta = \theta(\phi)$; $\phi \in \Phi$. This parameter is said to be *permissible* under the group G if

$$\theta(\phi_1) = \theta(\phi_2) \text{ implies } \theta(\phi_1 g) = \theta(\phi_2 g) \text{ for all } g \in G.$$

For a permissible parameter θ the transformations g can be defined in a unique way by $(\theta g)(\phi) = \theta(\phi g)$. In many simple cases the parameter θ will be permissible, but not always. It can be shown in general [9] that there always is a unique maximal subgroup G^0 of G such that the parameter θ is permissible under G^0 .

The *orbits* of a group G as acting upon a parameter space Θ is the set of parameter values obtained under the group by starting at a fixed parameter value θ_0 , i.e., the orbit in this case is given by $\{\theta_0 g; g \in G\}$. For a scalar parameter θ under the scale group $\theta \mapsto b\theta$; $b > 0$ there are three orbits: $\{\theta : \theta > 0\}$, $\{\theta : \theta < 0\}$ and then an orbit given by the single point 0. Both the groups

$\theta \mapsto \theta + a$ and $\theta \mapsto b\theta + a$ have one orbit consisting of the whole space Θ . In this case, when there is only one orbit, we say that the group action is *transitive*.

For a vector parameter θ , the group of rotations $\theta \mapsto C\theta$; $C^t C = I$, $\det(C) = 1$ has orbits consisting of the shells with constant $\|\theta\|$, while the group of all non-singular matrix multiplications: $\theta \mapsto A\theta$; $\det(A) \neq 0$ is nearly transitive; precisely, it has two orbits $\{\theta : \theta \neq 0\}$ and $\{0\}$.

There is a general statistical inference theory under symmetry, that is, for a statistical model endowed with a group G , for a summary, see [9]. One important result is the following: Within the orbits of a group acting upon the parameter space, one can always find an optimal estimator in a welldefined sense. This puts limitation upon the ways model reductions should be made, more precisely: In any statistical setting where a group is defined in a meaningful way, I will assume that every model reduction is to an orbit or to a set of orbits of the group acting upon the parameter space. Several examples motivating this policy further are given in [9].

I will finally need some concepts from group representation theory. A set of unitary operators $U(g); g \in G$ acting upon a Hilbert space $\mathbf{H}$ is called a representation of the group G if $U(g_1 g_2) = U(g_1)U(g_2)$ for all $g_1, g_2 \in G$, so that these operators form a new group, homomorphic to G .

A space $V \subset \mathbf{H}$ is invariant if $U(g)v \in V$ whenever $v \in V$ and $g \in G$. An invariant space is called irreducible if it does not contain any proper invariant subspace. As is well known, unitary representations, invariant spaces and irreducible spaces play an important rôle in many branches of quantum mechanics.

A simple and important example is the regular representation U_R , defined on the Hilbertspace $L^2(\Phi, \nu)$ of functions f on the total parameter space Φ by

$$U_R(g)f(\phi) = f(\phi g). \quad (2)$$

Assume now that θ is a permissible parameter. Then it is an easy exercise to show that

$$V = \{f : f(\phi) = \tilde{f}(\theta(\phi))\} \quad (3)$$

is an invariant space under U_R , and that

$$U_R(g)f(\phi) = \tilde{f}(\theta(\phi g)) = \tilde{f}((\theta g)(\phi)) \quad (4)$$

on V .

4 The new setting for quantum mechanics.

I am now ready to formulate what I consider to be the most natural set of axioms of quantum theory, axioms that may be motivated from the statistical discussion above. I will aim at deriving the Rules 1-3 of the Introduction from these axioms in the remaining parts of this paper, and in fact, I will derive more than that.

In contrast to the traditional set given by Rules 1-3, I consider most of the axioms below to be relatively natural in the light of common sense and in the

light of basic statistics in the way it has been formulated above. It is the totality of assumptions which makes the core of quantum theory in the way I see it.

Axiom 1. *For a given closed system there is total parameter space Φ whose elements $\phi \in \Phi$ are not estimable relative to any experient. There is a transformation group G defined on Φ . The space Φ is locally compact, and the group G is transitive on Φ . The right invariant measure under G on Φ is called ν .*

Axiom 2. *There is a set $\mathbf{A}$ of potential experiments $\mathbf{E}^a; a \in \mathbf{A}$ on this system. For each $a \in \mathbf{A}$ there corresponds a maximal parameter $\theta^a = \theta^a(\phi)$. The parameter θ^a may or may not be estimable.*

Axiom 3. *For the case where θ^a is not estimable, there is a model reduction $\lambda^a = \lambda^a(\theta^a)$ such that λ^a is estimable under the experiment $\mathbf{E}^a$. Assume that λ^a is maximal under this requirement.*

This means that the remaining part of θ^a , that is, fixing λ^a , is not estimable from any experiment.

Axiom 4. *For each $a \in \mathbf{A}$ there is a realization of the experiment $\mathbf{E}^a$ (that is, a measuring instrument) such that the observation y has a complete sufficient statistic t^a .*

Axiom 5. *When λ^a is discrete, Axiom 4 can be strengthened to the existence of a perfect experiment for λ^a , an experiment where measurement noise can be disregarded and where the resulting (pure) states are determined by statements of the form $\lambda^a = \lambda_k^a$.*

Given the result of a perfect experiment $\mathbf{E}^a$, we can predict the result of a new perfect experiment $\mathbf{E}^b$. When λ^a takes only two values, we must also assume that such a probabilistic prediction exists when we allow for a final state being a mixed state or an effect; for definitions and interpretations see later.

Definition 1. *Let G^a be the maximal subgroup of G with respect to which $\lambda^a(\phi)$ is a permissible function of ϕ . This group will always exist.*

For definiteness, regarding statistical methods, it is natural to assume Bayesian estimation in each experiment with a prior induced on each λ^a by ν on Φ . This is also equivalent to what is called the best equivariant estimator under the group G^a , see [9].

Axiom 6. *(i) For each pair of experiments $\mathbf{E}^a, \mathbf{E}^b; a, b \in \mathbf{A}$ there is an element k_{ab} of the basic group G which induces a correspondence between the respective reduced parameters:*

$$\lambda^b = \lambda^a k_{ab} \text{ or } \lambda^b(\phi) = \lambda^a(\phi k_{ab}). \quad (5)$$

(ii) The set $K = \{k_{ab}; a, b \in \mathbf{A}\}$ of transformations upon Φ constitutes a subgroup of G .

Using the fact that (i) implies that, given g^a , one must have $\lambda^a g^a$ equal to $\lambda^b g^b k_{ba}$ for some g^b , then using (i) again, it follows that $\lambda^a(\phi g^a) = \lambda^a(\phi k_{ab} g^b k_{ba})$ for all a . Hence, since G^a is determined by its actions on λ^a : $\lambda^a(\phi g^a) = (\lambda^a g^a)(\phi)$, it follows that $g^a = k_{ab} g^b k_{ba}$. What we really need later in this paper is the implied relation between unitary representations

$$U(g^a) = U(k_{ab})U(g^b)U^\dagger(k_{ab}). \quad (6)$$

The relation $g^a = k_{ab} g^b k_{ba}$ is what mathematicians call an inner homomorphism between group elements, or really an isomorphism. An isomorphism means that essentially the same group is acting upon both spaces Λ^a and Λ^b , and often in such cases the same group element symbol is used. We will use different symbols, however, because the actions are related to different experiments.

The symmetry assumptions between experiments given in Axiom 6 will turn out later to be crucial for the development of quantum theory.

Axiom 7. The groups $G^a; a \in \mathbf{A}$ generate G .

We will show later that these axioms leads to the existence of a Hilbert space $\mathbf{H}^a$ for the experiment $\mathbf{E}^a$ in such a way that there are basis vectors $f_k^a \in \mathbf{H}^a$ which can be uniquely coupled to the statements that a perfect measurement of λ^a gives the result that $\lambda^a = \lambda_k^a$ for suitable constants λ_k^a . These Hilbert spaces can be realized as subspaces of $L^2 = L^2(\Phi, \nu)$, where ν is the invariant measure under G assumed in Axiom 1. In particular, $\mathbf{H}^a$ is a subspace of

$$V^a = \{f \in L^2 : f(\phi) = \tilde{f}(\lambda^a(\phi))\}. \quad (7)$$

Both $\mathbf{H}^a$ and V^a are invariant under the regular representation of the group G^a ; in simple cases the two are equal. In general we have one of two cases: Either $\mathbf{H}^a$ is an irreducible space under G^a or we have a decomposition $\mathbf{H}^a = \mathbf{H}_1^a \oplus \mathbf{H}_2^a \oplus \dots$, where each $\mathbf{H}_i^a$ is irreducible under G^a .

For different a , the Hilbert spaces $\mathbf{H}^a$ turn out to be unitarily connected, and this is crucial. It implies that there exists a common Hilbert space $\mathbf{H}$ such that $\mathbf{H}^a = U^a \mathbf{H}$ for unitary U^a . The space $\mathbf{H}$ is an invariant space of the whole group G under the representation exemplified by

$$W(g_1^a g_2^b g_3^c) = U^{a\dagger} U(g_1^a) U^a U^{b\dagger} U(g_2^b) U^b U^{c\dagger} U(g_3^c) U^c \quad (8)$$

If $\mathbf{H}$ is not irreducible under $\{W(g)\}$, then we have a decomposition $\mathbf{H} = \mathbf{H}_1 \oplus \mathbf{H}_2 \oplus \dots$. This decomposition will correspond to the well known superselection rules of quantum mechanics.

The last question is then if all normalized vectors in each component $\mathbf{H}_i$ can be considered as a state vector. This will be proved later to follow from the other assumptions above together with the following axiom, which may be expected to hold if there are enough potential experiments involved.

Axiom 8. *The unitary group generated by $\{W(g)\}$ and the phase factors $e^{i\alpha}$ is transitive on each component $\mathbf{H}_i$.*

5 A large scale model.

One of the simplest non-commutative groups is the group S_3 of permutations of 3 objects. It has a two-dimensional representation discussed in many books in group theory and in several books in quantum theory. The quantum theory book by Wolbarst [16] is largely based upon this group as a pedagogical example.

I will in this Section visualize this group by considering the permutations of the corners of an equilateral triangle, which can be realized physically by the change of position of some solid version of this triangle. This will serve to illustrate the axioms above on a macroscopic example.

The spatial orientation ϕ of the whole triangle will be looked upon as a hidden total parameter, and to this end we will place the solid triangle within a hollow nontransparent sphere, with the corners on the sphere, in such a way that it can rotate freely around its center point, placed at the center of the sphere. The basic group G is to begin with taken as the group of such rotations, but later, when we specialize to the corners, we will take G as the permutation group. Let the solid triangle be painted white on one side and black on the other side.

Let there be 4 small windows in the sphere, one at the north pole, where the colour facing up can be observed, and three equidistant windows along the equator, where the closest corner of the triangle can be observed.

The measurements made in the windows could be uncertain for some reason, and we could model this in the ordinary statistical way by some model $Q^\lambda(dy)$, depending upon a (reduced) parameter which can be thought about as the ideal measurement.

Hence there are 4 reduced parameters, corresponding to the 4 different experiments that can be done in this case, one for each window: λ^0 is the ideal colour as observed from the north pole window: λ^a , λ^b and λ^c are the three ‘correct’ corners of the triangle as observed from the windows a , b and c , respectively. The term ‘correct’ will be defined more precisely below. The parameter λ^0 takes the values Bl and Wh , and the parameters λ^i for $i = a, b, c$ each takes the values A , B and C , say. All these parameters can be considered to be functions of the triangle’s spatial orientation ϕ within the sphere.

Lemma 1 *Both with respect to the group of permutations and with respect to the group of rotations, λ^0 is a permissible function, while λ^a , λ^b and λ^c each are non-permissible. The largest permutation group with respect to which λ^a is permissible, is the group of cyclic permutations of the corners of the triangle, similarly for λ^b and λ^c .*

Proof.

Consider the 6 elements of the group S_3 of permutations: $g_1(ABC \mapsto ABC)$, $g_2(ABC \mapsto CAB)$, $g_3(ABC \mapsto BCA)$, $g_4(ABC \mapsto ACB)$, $g_5(ABC \mapsto CBA)$, $g_6(ABC \mapsto BAC)$.

Assume $\lambda^0(\phi_1) = \lambda^0(\phi_2)$, say black, for two total parameter values ϕ_1 and ϕ_2 . Then by simple inspection, $\lambda^0(g_i\phi_1) = \lambda^0(g_i\phi_2) = Bl$ for $i = 1, 2, 3$ and Wh for $i = 4, 5, 6$. Hence λ^0 is permissible.

For the other functions it is enough to produce a counterexample. Here is one for λ^a : Let ϕ_1 be any hyperparameter value, and by definition let ABC be the sequence of corners in ϕ_1 corresponding to the windows a, b, c . Put $\phi_i = \phi_1 g_i$ for $i = 2, \dots, 6$. Then $\lambda^a(\phi_1) = \lambda^a(\phi_4) = A$, but $\lambda^a(\phi_1 g_5) = \lambda^a(\phi_5) = C$ and $\lambda^a(\phi_4 g_5) = \lambda^a(\phi_3) = B$. Since the group elements g_4 and g_6 have the same structure as g_5 : Permutation fixing one corner of the triangle, a similar statement holds for these group elements.

To check that λ^a is permissible under the cyclic group, we can use direct verification. The details are omitted. A geometric proof is simpler than an algebraic proof.

One can easily imagine that an ideal measurement at the window a in principle can give more information θ^a about the position of the triangle, but that this information is hidden. One way to make this precise, is the following: Let us divide the sphere into 3 sectors corresponding to each of the windows a, b and c by using the meridians midway between a and b , midway between a and c and midway between b and c as borders between the sectors. Let S^a be the sector containing window a . Define θ^a as 1) the points among the triangle corners A, B and C that happen to be in the sector S^a ; 2) the coordinates of any two points which happen to belong to the same sector. By a geometric consideration, it can be seen that θ^a is permissible with respect to the subgroup G^a of rotations around the polar axis, and only permissible with respect to this subgroup.

From the geometry, it can be seen that S^a can contain 0, 1 or 2 triangle corners. This can be used to define λ^a precisely: If S^a contains 1 corner, let this corner be λ^a . If S^a contains 2 corners, let the closest one, as calculated from the coordinates, be λ_a . If S^a contains 0 corners, then exactly one of its neighbouring sectors must contain 2 corners. One is then chosen to be λ^b , respectively λ^c ; let the other one be λ^a . Since the coordinates of the corners that are in the same sector are contained in θ^a , it is seen that λ^a is a function of θ^a .

Note that the reduction from ϕ via θ to the parameter λ is forced upon us in this situation by the limitation in the possibility to make observations on the system.

We assume that there is some mechanism to ensure that it is impossible to look through two equatorial windows at the same time.

The Axioms 1, 2 and 3 are easily seen to be satisfied for this example. As for the Axiom 5 and the first part of axiom 5, they can be assumed to hold via some observational system. As for the possibility of prediction in the last part of Axiom 5, this follows from the symmetry of the system. Annihilation and creation is not relevant for this example. Again, Axioms 6 is easily satisfied.

Axiom 7 is not satisfied. The basic group S_3 has an irreducible two-dimensional representation. Then, from the theory which we will develop later, it is possible to predict probabilistically, from an observation in window a , the results of an observation in window b immediately afterwards.

However, in this case the prediction will be rather trivial, something that can be related to the fact that the subgroup under which the observation in a window is permissible, is the same for all windows. More complicated examples using the same concept can easily be imagined.

6 The spin 1/2 particle.

The most simple quantum mechanical system, a qubit, is realized as an electron with its spin. The spin component λ can be measured in any spatial direction a , and λ always takes one of the values -1 and +1.

In this section, we will give a non-standard, but quite intuitive description of a particle with spin, a description which we later will show to be equivalent to the one given by ordinary quantum theory.

Look first at a general classical angular momentum. A total parameter ϕ corresponding to such an angular momentum may be defined as a vector in three dimensional space; the direction of the vector giving the spin axis, the norm gives the spinning speed. A possible associated group G is then the group of all rotations of this vector in $\mathbf{R}^3$ around the origin together with the changes of the norm of the vector.

Now let the electron at the outset have such a total parameter ϕ attached to it, and let $\kappa = \|\phi\|$. It is of course well known that it is impossible to obtain in any way such detailed information about the electron spin. Now let us forget for a moment what we know about the electron, and assume that we set forth in an experiment $\mathbf{E}^a$ to measure its angular momentum component $\theta^a(\phi) = \kappa \cos(\alpha)$ in some direction given by a unit vector a , where α is the angle between ϕ and a . The measurement can be thought of as being done with a Stern-Gerlach device, which strictly speaking measures an observable y whose distribution depends upon θ^a , implying a possibility that the parameter θ^a - or some part of it - can be estimated from such a measurement. Given a , and given the measurement in the direction a , the rest of the total parameter ϕ will be unestimable.

With respect to the group G , the function $\theta^a(\cdot)$ is easily seen to be non-permissible for fixed a , simply because two vectors with the same component along a in general will have different such components after a rotation. The maximal possible group G^a with respect to which θ^a is permissible, is the group generated by the rotations of the vector ϕ around the axis a possibly together with a 180° rotation around any axis perpendicular to a , plus a possible scale change $\kappa \mapsto b\kappa; b > 0$.

In analogy to the situation in the previous Section, assume now that the electron's total parameter ϕ always is hidden, in such a way that for every a , the only part of $\kappa \cos(\alpha)$ we are able to measure, is the value +1 or -1, giving the sign of this component. We call this part $\lambda^a(\phi)$. This is an extreme model

reduction compared to θ^a , but interestingly enough, the model reduction is to an orbit of the group G^a .

The measured part found by the Stern-Gerlach apparatus, which may contain additional measurement noise, also takes the values ± 1 , and is called $\hat{\lambda}^a$. In some instances below, we will disregard such measurement noise, and assume the ideal condition $\hat{\lambda}^a = \lambda^a$. Such an approximation makes sense, also from a statistical point of view, for a discrete parameter.

Finally, since the model reduction is to a parameter of fixed norm, we delete the scale change part from the groups G and G^a . In particular, G is the group of rotations.

The Axioms formulated earlier can again be verified for the electron as described above. The electron as such has a Hilbert space with only one component, but for some given particle which can have spin 0, $1/2$, 1, $3/2$ or $\dots$, the general representation of the rotation group gives a decomposition as indicated.

7 Hilbert space of a single experiment.

In the following sections we will derive the usual quantummechanical description of the electron from the simple description above, including the Hilbert space with the associated interpretation of state vectors, the transition probabilities found from Born's rule and the representation of the state vectors for the electron spin on the Bloch or Poincaré sphere. An important point, however, is that we will make the derivations in general terms, and to this end we will take our basic Axioms as a point of departure. Because of space limitation, the derivations will be rather brief, but we hope to expand on this elsewhere.

We first look at the space S of observations, which we for simplicity can assume is common for all potential experiments. By Axiom 4 we assume that for each experiment $\mathbf{E}^a$ there exists a complete sufficient statistics t^a under the (reduced) model Q^{λ^a} for this experiment. Through the statistical model, a transformation group on the sample space of an experiment induces a transformation group on the parameter space. In [9] it is shown that for a complete, sufficient statistics one can also go the other way. Thus G^a (Definition 1) defines a transformation group, also called G^a , on S . From the Axiom 7 it follows that the groups G^a together generate the full group G , which then can be assumed defined on S and having a right invariant measure Q on that space. Without loss of generality we can - and will - assume that all the model measures Q^{λ^a} are absolutely continuous with respect to Q . All this must be considered to be details with the main purpose of establishing the measure Q on S .

We now define the Hilbert space $\mathbf{K}^a$ as the set of all complexvalued functions $h(t^a) \in L^2(S, Q)$ such that $f(\phi) = E^{\lambda^a}(h(t^a)) \in L^2(\Phi, \nu)$. Since sufficiency and completeness are properties that are invariant under group transformations, it follows that $\mathbf{K}^a$ is an invariant space for the regular representation of the group G on $L^2(S, Q)$.

Define now the operator A^a from $\mathbf{K}^a$ to $V^a = \{f \in L^2(\Phi, \nu) : f(\phi) =$

$$\tilde{f}(\lambda^a(\phi)) \text{ by} \quad (A^a y)(\lambda^a(\phi)) = E^{\lambda^a}(y) \quad (9)$$

Definition 2. Define the space $\mathbf{H}^a \subset V^a$ by $\mathbf{H}^a = A^a \mathbf{K}^a$.

It is easy to see that $\mathbf{H}^a$ is a closed subspace of $L^2(\Phi, \nu)$, and therefore a Hilbert space.

Proposition 1. The space $\mathbf{H}^a$ is an invariant space for the regular representation of the group G^a .

A main result is now:

Theorem 1. The spaces $\mathbf{K}^a$ and $\mathbf{H}^a$ are unitarily related. Also, the regular representations of the group G^a on these spaces are unitarily related.

To prove this, one has to show that the mapping A^a can be replaced by a unitary map in the relation $\mathbf{H}^a = A^a \mathbf{K}^a$. This is shown in detail in [10] by first constructing an explicit relation between the two regular representations and then using a general mathematical result stating that two representations are unitarily equivalent if they are equivalent.

8 The quantum theoretical Hilbert space.

The regular representation U restricted to the subgroup G^a can also act upon functions of λ^a , similarly for the representation restricted to G^b . Then both $\mathbf{H}^a$ and $\mathbf{H}^b$ are invariant spaces for these respective representations by Proposition 1. In this Section we will assume that these invariant spaces are irreducible; the general case is treated in later. We need

Schur's lemma [2] *If U_1 and U_2 are irreducible representations on Hilbert spaces $\mathbf{H}_1$ and $\mathbf{H}_2$, and A is such that $U_1(g_1)A = AU_2(g_2)$ for all g_1 and some corresponding g_2 , then either $A = 0$ or A is an isomorphism between the spaces $\mathbf{H}_1$ and $\mathbf{H}_2$.*

(In [2] one assumes $g_1 = g_2$, which is related to the tradition in pure mathematics to take abstract groups as points of departure. By contrast, we talk about transformation groups, and then it may be natural sometimes to use different symbols for transformations on different spaces.)

Recall from Axiom 6 and (6) that $U(g^a)U(k_{ab}) = U(k_{ab})U(g^b)$. From this and Schur's lemma on each irreducible component it follows that we can construct a connection between the spaces by

$$\mathbf{H}^b = U(k_{ab})\mathbf{H}^a \quad (10)$$

Then (10) and Theorem 1 gives an important result:

Theorem 2. *There is a Hilbert space $\mathbf{H}$, and for each $a \in \mathbf{A}$ a unitary transformation U^a such that $\mathbf{H}^a = U^a \mathbf{H}$. There are also unitary transformations V^a such that the observational Hilbert spaces satisfy $\mathbf{K}^a = V^a \mathbf{H}$.*

By Axiom 7, $\{G^a; a \in \mathbf{A}\}$ generate the whole group G . From this it follows:

Theorem 3. *The basic Hilbert space $\mathbf{H}$ is an invariant space for the whole group G .*

Proof: From Proposition 1, $\mathbf{H}^a$ is an invariant space for the group G^a under the regular representation U . Assume now that $g = g_1 g_2 g_3$, where $g_1 \in G^a$, $g_2 \in G^b$ and $g_3 \in G^c$. Then

$$W(g_1 g_2 g_3) = U^{a\dagger} U(g_1) U^a U^{b\dagger} U(g_2) U^b U^{c\dagger} U(g_3) U^c \quad (11)$$

gives a representation on $\mathbf{H}$ of the set of elements of G that can be written as a product $g_1 g_2 g_3$ with $g_1 \in G^a$, $g_2 \in G^b$ and $g_3 \in G^c$. Continuing in this way we are able to construct a representation of G on the space $\mathbf{H}$.

9 The finitedimensional case with perfect measurements.

The discussion up to now has been very general, but in the rest of this paper I will concentrate on the case where $\mathbf{H}$ has a finite dimension n . Then by the unitary connection just proved, each of the spaces $\mathbf{H}^a$ will also be n -dimensional.

Now $\mathbf{H}^a$ is a function of $\lambda^a = \lambda^a(\phi)$. Assume for simplicity that λ^a is a scalar; since it can be assumed to take a finite number of values, this can always be arranged for. Define on this space an operator S^a by

$$S^a f(\lambda^a) = \lambda^a f(\lambda^a). \quad (12)$$

As a Hermitian operator on an n -dimensional Hilbert space, this will have n eigenvalues λ_k^a and corresponding eigenfunctions f_k^a .

Using Axiom 5, we will from now on assume that each experiment $\mathbf{E}^a$ is perfect, that is, measurement error can be disregarded, so that $\hat{\lambda}^a = \lambda^a$. Then we will show that the eigenfunctions above can be chosen in a particularly simple way, there is no degeneracy, and $\mathbf{H}^a$ will be the whole space $V^a = \{f \in L^2 : f(\phi) = \tilde{f}(\lambda^a(\phi))\}$.

Theorem 4. *(i) For a perfect experiment $\mathbf{E}^a$ associated with a finite-dimensional Hilbert space, the space $\mathbf{H}^a$ contains the normalized functions $f_k^a(\phi) = \sqrt{n} I(\lambda^a(\phi) = \lambda_k^a)$.*

(ii) These functions are eigenfunctions for S^a corresponding to eigenvalues λ_k^a . There is no degeneracy in this case.

(iii) We can take $\mathbf{H}^a$ to be equal to V^a in this case.

Proof: (i) Since the experiment is perfect and by Axiom 3 has a complete sufficient statistic t^a which then [13] is minimal, this t^a must take a finite number of values t_k^a in such a way that t_k^a leads to the certain conclusion $\lambda^a = \lambda_k^a$. Let $h_k^a(t^a) = \sqrt{n}I(t^a = t_k^a)$, and define $f_k^a(\phi) = E^{\lambda^a} h_k^a(t^a) = P^{\lambda^a}(t^a = t_k^a)$. Since the posterior probability in general is proportional to the likelihood times the prior, and the prior in this case is uniform on the discrete λ -values, this must imply that the likelihood is $f_k^a(\phi) = \sqrt{n}I(\lambda^a(\phi) = \lambda_k^a)$. By definition, these must belong to $\mathbf{H}^a$.

(ii) The first part is obvious. For these particular eigenfunctions, equal eigenvalues implies equal eigenfunctions, so there is no degeneracy.

(iii) The fact that $\mathbf{H}^a$ is n -dimensional, means that the n eigenfunctions $f_k^a(\phi) = \sqrt{n}I(\lambda^a(\phi) = \lambda_k^a)$ span $\mathbf{H}^a$ for the n different eigenvalues λ_k^a . But these functions also span the total set of functions on $\lambda_1^a, \dots, \lambda_n^a$, i.e., V^a .

Remark: Using the L^2 -norm in V^a , the functions f_k^a above are normalized, since the invariant (probability) measure has mass $1/n$ on each λ_k^a .

Fix c such that $\mathbf{H} = \mathbf{H}^c$, which has a basis $\{f_j^c\}$ given by $f_j^c(\phi) = \sqrt{n}I(\lambda^c(\phi) = \lambda_j^c)$. In a different space $\mathbf{H}^a$ these correspond to $f_j^a(\phi) = U(k_{ca})f_j^c(\phi)$. Now define vectors

$$v_j^a = W(k_{ca})f_j^c, \quad (13)$$

where W is the representation (11). These are eigenvectors of the selfadjoint operator

$$T^a = W(k_{ca})S^aW(k_{ac}) \quad (14)$$

with eigenvalues $\lambda_j^a = \lambda_j^c k_{ca}$.

The information contained in the vector v_k^a is simply: The perfect experiment $\mathbf{E}^a$ has been performed, and the result is that $\lambda^a = \lambda_k^a$.

The operator T^a may be written

$$T^a = \sum_1^n \lambda_k^a v_k^a v_k^{a\dagger}. \quad (15)$$

These operators are selfadjoint.

Remark: The present operator S^a and thus T^a has a non-degenerate spectrum. This corresponds to the fact that the parameter λ^a - though it has also been called a reduced parameter relative to a parameter θ^a which do not take a value - is a full parameter for the experiment $\mathbf{E}^a$. To achieve degenerate spectra, one must look upon partial parameters $\mu^a = \mu^a(\lambda^a)$, i.e., spin of one particle in a two-particle system or the mass as one of several quantum numbers associated with a particle.

10 Invariance, states and superselection rules.

In the previous Section we started with indicator functions f_k^a in $\mathbf{H}^a$ as eigenfunctions of S^a with eigenvalues λ_k^a . These are transformed in the natural way by the group elements $g^a \in G^a$ and by the corresponding regular unitary representation operators.

Theorem 5. *If f_k^a is an eigenvector for S^a with eigenvalue λ_k^a , then $U(g^a)f_k^a$ is an eigenvector with eigenvalue $\lambda_k^a g^a$.*

The proof uses the fact that $U(g^a)f_k^a(\phi) = \tilde{f}_k^a((\lambda^a g^a)(\phi))$, so $S^a f_k^a = \lambda_k^a f_k^a$ implies $U^\dagger(g^a)S^a U(g^a)f_k^a(\phi) = \lambda_k^a g^a f_k^a(\phi)$.

Corollary 1. *Let $W(g^a) = U^{a\dagger}U(g^a)U^a$ be the operator on the basic Hilbert space $\mathbf{H}$ corresponding to $U(g^a)$ on $\mathbf{H}^a$. Then the following holds: If v_k^a is an eigenvector for T^a with eigenvalue λ_k^a , then $W(g^a)v_k^a$ is an eigenvector with eigenvalue $\lambda_k^a g^a$.*

It is a general theorem from representation theory [2] that every finitedimensional invariant space can be decomposed into irreducible invariant spaces. Thus we have $\mathbf{H}^a = \mathbf{H}_1^a \oplus \mathbf{H}_2^a \oplus \dots$, where the $\mathbf{H}_i^a$ are irreducible.

By doing this in both the experiments $\mathbf{E}^a$ and $\mathbf{E}^b$ we see that we can assume that the relation $U(g^a)U(k_{ab}) = U(k_{ab})U(g^b)$ holds separately between irreducible spaces on each side. From this, Schur's lemma gives that we have relations of the form $\mathbf{H}_i^a = U(k_{abij})\mathbf{H}_j^b$ between all these spaces, or else $U(k_{abij}) = 0$ with no relation at all. This gives unitary relations between pairs of irreducible spaces, one for each experiment, and the space $\mathbf{H}^b$ has a conformable decomposition $\mathbf{H}_1^b \oplus \mathbf{H}_2^b \oplus \dots$.

It follows also that $\mathbf{H} = \mathbf{H}_1 \oplus \mathbf{H}_2 \oplus \dots$ conformably, and that there is a unitary connection $\mathbf{H}_i^a = U_i^a \mathbf{H}_i$.

Summarizing the above discussion we have:

Theorem 6. *The basic Hilbert space $\mathbf{H}$ can be decomposed as $\mathbf{H} = \mathbf{H}_1 \oplus \mathbf{H}_2 \oplus \dots$, where each $\mathbf{H}_i$ is an irreducible invariant space under the group G . All unit vectors in the Hilbert space under this decomposition are possible state vectors. Each part corresponds to a fixed value of one or several quantities that are conserved under all experiments.*

It remains to prove that all $v \in \mathbf{H}_i$ are possible state vectors under Axiom 8. To this end, first observe that all state vectors are of the form $v_k^a = W(g^c k_{ca})f_j^c$ for some $g^c \in G^c$ and $k_{ac} \in K$.

Now every group element $g \in G$ is of the form $g^a k_{ab}$ for some b , some $g^a \in G^a$ and some $k_{ab} \in K$. For instance, let $g = g_1^b g_2^c g_3^d$ for $g_1^b \in G^b$, $g_2^c \in G^c$ and $g_3^d \in G^d$. Then

$$g = g_1^b g_2^c (g_3^c k_{cd}) = g_1^b g_4^c k_{cd} = g_1^b (g_4^b k_{bc}) k_{cd} = g_5^b k_{bd} = g_5^a k_{ab} k_{bd} = g_5^a k_{ad}.$$

From this, the set of state vectors coincides with the set of vectors of the form $W(g)v_0$ for some $v_0 = f_j^c$.

Therefore, under Axiom 8 $\{U(g)v_k^a; g \in G\}$ consists of all unit vectors in $\mathbf{H}_i$, and all unit vectors in each irreducible space $\mathbf{H}_i$ are state vectors.

An alternative way to formulate a non-trivial decomposition $\mathbf{H} = \mathbf{H}_1 \oplus \mathbf{H}_2 \oplus \dots$ may be to say that not all linear combinations of state vectors are state vectors, in particular not linear combinations of vectors that correspond to different values of absolutely conserved quantities like mass or charge. These superselection rules are introduced in an ad hoc way in standard quantum mechanics, and they are neglected in many textbooks. They arrive in a quite natural way in our approach.

11 The Born formula.

So far I have constructed the quantummechanical Hilbert space from my setting, and I have given a simple interpretation of the vectors of this Hilbert space: They can all be interpreted in a question-and answer form: 1) Choose a and ask the question: What is the value of λ^a ? 2) Do a (perfect) experiment and give the answer: $\lambda^a = \lambda_k^a$.

What is left, is to associate probabilities to these states. In this paper I follow Cartwright [4] and say that all probabilities come from transition probabilities. These can be found from Born's formula. Later I will show, in a similar way as other authors have done it, that the ordinary quantum formalism follows from this.

In this and the next Section I will concentrate on the case with one irreducible component in the Hilbert space, i.e., I will neglect superselection rules. This is really no restriction, since transitions always are within one component.

By Axiom 5 there exist transition probabilities in the sense that if we know that $\lambda^a = \lambda_k^a$, there is a well-defined probability that a later perfect experiment $\mathbf{E}^b$ will result in $\lambda^b = \lambda_i^b$. We must first investigate what it means that two such final states correspond to the same state vector v .

Lemma 2. *Consider a system where each experimental parameter λ only takes two values: -1 and +1. Assume that two vectors in $\mathbf{H}$ satisfy $v_i^b = v_j^c$, where v_i^b corresponds to $\lambda^b = \lambda_i^b$ for a perfect experiment $\mathbf{E}^b$ and v_j^c corresponds to $\lambda^c = \lambda_j^c$ for a perfect experiment $\mathbf{E}^c$. Then there is a 1-1 function F such that $\lambda^c = F(\lambda^b)$ and $\lambda_j^c = F(\lambda_i^b)$.*

Proof: The assumption is that

$$U^{b\dagger} f_i^b = U^{c\dagger} f_j^c. \quad (16)$$

Without loss of generality, let both λ_i^b and λ_j^c be equal to 1. Then

$$\sqrt{n}I(\lambda^c(\phi) = 1) = f_j^c(\phi) = U^c U^{b\dagger} f_i^b(\phi) = \sqrt{n}I((\lambda^b k_{bc})(\phi) = 1).$$

Thus $I(\lambda^c = 1) = I(\lambda^b k_{bc} = 1)$. But this means that the two level curves coincide, and we must have $\lambda^c = \lambda^b k_{bc} = F(\lambda^b)$.

My aim in this Section is to prove Born's celebrated formula for transition probabilities in quantum mechanics. This result is related to the well known [7, 12] Gleason's theorem, and will be proved from this theorem (together with Lemma 2 above) in [10] for the case where the dimension of the Hilbert space is larger than 2.

In this paper I will concentrate on the case of dimension 2. Then it seems necessary to use stronger assumptions, as stated in Axiom 5 above. For this case I will also use a more suitable version of Gleason's theorem recently proved by Busch [3] and Caves et al [5], which also is valid for dimension 2.

Gleason's theorem variant. *Consider the set of effects on a Hilbert space $\mathbf{H}$, defined as the set of Hermitian operators having eigenvalues in the unit interval. Assume that there is a generalized probability measure π on these effects satisfying*

$$0 \leq \pi(E) \leq 1 \text{ for all } E.$$

$$\pi(I) = 1.$$

$$\pi(\sum E_j) = \sum \pi(E_j) \text{ whenever } \sum E_j \leq I.$$

Then there exists a density operator ρ , that is, a positive operator with unit trace, such that $\pi(E) = \text{tr}(\rho E)$.

Associated with this I now prove Born's formula:

Theorem 7. *Assume that we know that $\lambda^a = \lambda_i^a$ and that we are interested in the probability that a new perfect experiment $\mathbf{E}^b$ results in $\lambda^b = \lambda_j^b$. Then this transition probability is given by*

$$P(\lambda^b = \lambda_j^b | \lambda^a = \lambda_i^a) = |v_i^{a\dagger} v_j^b|^2. \quad (17)$$

As already stated, I will carry out the proof below only for dimension 2. It can be seen from the proof that I really show more than what is stated in Theorem 7: There is also a corresponding formula for transition to and from mixed states which follows. Furthermore, in the next Section I also discuss concrete interpretations of the rather abstract 'effects' for the case of dimension 2, and I actually here give a proof of the Gleason theorem variant for this case.

Proof: For this proof fix a and i and hence the state v_i^a , interpreted as $\lambda^a = \lambda_i^a$. Without loss of generality we can take $i = 1$ and $\lambda_i^a = 1$. By Lemma 2, every end state corresponding to the same vector $v = v_j^b$ must lead to the same transition probability. By Axiom 5 we can assume an extension of this class of transition probabilities, in the next Section shown to be equivalent to the transition to an arbitrary effect as defined above.

A general twodimensional Hermitian operator can be written $E = \frac{1}{2}(rI + cu \cdot \sigma)$, where r and c are scalar, u is a 3-vector of unit norm and σ is a vector with the 3 Pauli spin matrices as components:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The eigenvalues of E are $\frac{1}{2}(r \pm c)$. The factor $\frac{1}{2}$ is included for convenience.

In particular this representation holds for an effect, for which the conditions $0 \leq c \leq 1$ and $c \leq r \leq 2 - c$ ensure that the eigenvalues are between 0 and 1. A very particular case is the Bloch sphere or Poincaré sphere of a pure state density matrix $vv^\dagger$, which corresponds to the case $c = 1$ and $r = 1$. An intermediate case is a general density matrix with $r = 1$ and $c \leq 1$.

In this proof, and also in the next Section, we will study the set in the (r, c) -plane implied by the set of effects. Here the effects are confined to the interior of the triangle with corners $(0, 0)$ (giving $E = 0$), $(2, 0)$ (giving $E = I$) and $(1, 1)$ (corresponding to points on the Bloch sphere).

I start by assuming that there exists a transition probability $\pi(r, cu)$ from the fixed state v_1^a given by $\lambda^a = 1$ to an arbitrary effect $E = \frac{1}{2}(rI + cu \cdot \sigma)$. This will be interpreted concretely in the next Section.

The basic requirement given to probabilities on these effects is that $\sum E_i \leq I$ should imply $\sum \pi(r_i, c_i u_i) = \pi(\sum r_i, \sum c_i u_i)$. Arguments for this requirement from simple, general assumptions will be given in the next Section.

One way to satisfy this requirement, is to let $\pi(r, cu) = \frac{1}{2}(r + ck \cdot u)$ for some 3-vector k . We will start by studying this particular solution. First, we must have $\|k\| \leq 1$ in order to get a probability π in the interval $[0, 1]$ for pure states. The initial condition is that we shall have $\pi = 1$ when $r = 1$, $c = 1$ and $u = a$ with norm 1 corresponds to the pure state

$$\frac{1}{2}(I + a \cdot \sigma) = v_1^a v_1^{a\dagger}.$$

This gives $k \cdot a = 1$, and since a and k both have norm less than or equal to 1, then by necessity $k = a$.

But in fact, $\pi(r, cu) = \frac{1}{2}(r + ca \cdot u)$ is the only solution of the problem. This follows from the following result, proved by Caves et al. [5]: πE must be a linear function: $\pi(\alpha E) = \alpha \pi(E)$. Since $E = \frac{1}{2}(rI + cu \cdot \sigma)$ depends linearly upon 4 parameters r, cu_1, cu_2, cu_3 , it follows that $\pi(r, cu)$ must have the form $\beta r + \gamma \cdot (cu)$. Since $\pi(2, 0) = 1$, we must have $\beta = 1/2$. As just shown, this implies $\gamma = k/2 = a/2$.

So it remains to reformulate $\pi(r, cu) = \frac{1}{2}(r + ca \cdot u)$ for the case where the final effect is a pure state, i.e., $r = c = 1$. To this end, assume $vv^\dagger = \frac{1}{2}(I + u \cdot \sigma)$ and $v_1^a v_1^{a\dagger} = \frac{1}{2}(I + a \cdot \sigma)$. Then

$$\begin{aligned} |v_1^{a\dagger} v|^2 &= \text{tr}(v_1^a v_1^{a\dagger} vv^\dagger) = \text{tr}\left(\frac{1}{4}(I + a \cdot \sigma)(I + u \cdot \sigma)\right) \\ &= \text{tr}\left(\frac{1}{4}(I + (a + u) \cdot \sigma + a \cdot u I)\right) = \frac{1}{2}(1 + a \cdot u). \end{aligned}$$

Thus it follows that $\pi(r, u) = |v_1^{a\dagger} v|^2$ for this case.

Corollary 2. *If the initial state is given by a density function ρ_0 , then the transition probability to a final effect E is given by*

$$P(E|\rho_0) = \text{tr}(\rho_0 E).$$

Proof: An extension of the same proof. The basic point is that linearity must hold. Start with pure initial states and then take ρ_0 as a probability mixture.

The case of density matrices can be interpreted by standard quantum mechanics. The general case will be interpreted in the next Section.

12 Implications for the spin 1/2 particle and for simple decision problems.

12.1 A perfect measurement.

The point of departure in this Section is again that a state is given by some question plus the answer to that question, in short, the perfect measurement $\mathbf{E}^b$ together with $\lambda^b = \lambda_k^b$, or even shorter, just $\lambda^b = \lambda_k^b$. In the spin 1/2 case this will correspond to a choosen direction b , and the question about what the spin component in that direction is. The answer can be +1, corresponding in ordinary quantum mechanical terms to a certain eigenvector v_1^b , or -1, corresponding to the orthogonal eigenvector v_2^b .

But this means that there is a one-to-one correspondence between the possible state functions v (disregarding an irrelevant phase factor) and 3-vectors u : Define $u = b$ if a measurement in the direction b gives the result +1, and $u = -b$ if a measurement in the direction b gives the result -1. This is consistent: a measurement along $-b$ gives the value +1 if and only if a measurement along b gives the value -1.

Many textbooks discuss the Bloch sphere representation of this result:

$$v_1^b v_1^{b\dagger} = E_1(b) = \frac{1}{2}(I + b \cdot \sigma) \quad (18)$$

$$v_2^b v_2^{b\dagger} = E_2(b) = \frac{1}{2}(I - b \cdot \sigma) \quad (19)$$

I will let the matrix $E(u) = E_1(u)$ represent the statement that a perfect measurement along the direction a has given the value +1 when $u = b$, and the value -1 when $u = -b$. This is again consistent, and $\{E(u)\}$ is in one-to-one correspondence to the unit 3-vectors $\{u\}$.

This gives the specification of a state corresponding to a perfect measurement, and every quantummechanical state vector for a spin 1/2 particle can be

represented in this way: A question about a spin component in some direction, and the answer from a perfect experiment. This is perfectly in agreement with the general results given in earlier sections of this paper; for qubits we can, as just shown, device a more direct construction.

The issue of transition probabilities comes after this in my treatment. Assume some initial direction a . The state corresponding to the result that a measurement in this direction has resulted in $+1$, can be called $v(a)$ or $E(a)$ or simply a . Starting from this, one can compute the probability of getting $+1$ in a new direction b by Born's formula. The result can be expressed in many ways, a simple one being $P(b|a) = (1 + \cos(w))/2$, where w is the angle between a and b .

This is a derived result from Born's formula and the theory related to the state characterization. The characterization of the state itself can most simply in my view be given as above.

Born's formula is again, in the view of the present paper, a result of the assumption that the probabilities exist, not only from pure states to pure states, but to mixed states and also to the more general states called effects. Thus it becomes important to study these extended state concepts from a statistical point of view.

12.2 Allowing measurement errors.

Let the Stern-Gerlach apparatus be imperfect: Given that the real value is $+1$, it gives -1 with a certain probability and given that the real value is -1 , it gives $+1$ with some probability. Then a measurement $+1$ can be obtained in two ways: The right way or the wrong way.

From this model, from data and from some prior on the two states, one finds posterior probabilities for the two states,

Assume then that the reported state is $+1$, and that there from this is an aposteriori probability $p_1 \leq 1/2$ that the correct state is -1 . Then the true state is found by probability weighing:

$$E_1(b, p_1) = (1 - p_1) \frac{1}{2}(I + b \cdot \sigma) + p_1 \frac{1}{2}(I - b \cdot \sigma) = \frac{1}{2}(I + (1 - 2p_1)b \cdot \sigma) \quad (20)$$

It is well known from the theory of Bloch spheres that any mixed state can be found by replacing b for the pure state by cb for $0 \leq c < 1$. Here $1 - 2p_1$ is a general c in this interval. Hence any mixed state can be represented by such a measurement, in particular, by a vector b and an error probability p_1 .

An important point is that both p_1 and the unit vector b can be recovered if the density matrix E_1 is given.

Similarly, if the reported state is -1 , and there is an aposteriori probability $p_2 \leq 1/2$ that the correct value is $+1$, one gets the true state

$$E_2(b, p_2) = (1 - p_2) \frac{1}{2}(I - b \cdot \sigma) + p_2 \frac{1}{2}(I + b \cdot \sigma) = \frac{1}{2}(I - (1 - 2p_2)b \cdot \sigma) \quad (21)$$

Of course a similar state can be found with prior probabilities instead of posterior probabilities, say with a prior π for $+1$. The very common case $\pi = 1/2$ gives a totally uninformative mixed state $\frac{1}{2}I$.

The transition probability to other pure or mixed states can be found from Corollary 2.

12.3 Information on both types of errors.

The calculation in the previous subsection was for the case where p_1 and p_2 were posterior probabilities for the two types of errors. But a similar calculation applies for the case where p_1 and p_2 are replaced by predetermined error probabilities as in statistical testing of hypotheses. For the classical theory of hypothesis testing, any statistical textbook can be consulted.

So we will look upon a hypothesis testing problem with level α and power β . In our connection this means the following: Before any data are obtained, we make a programme stating how our decision procedure shall be. This goes as follows: The decision shall be based upon an observator t^a which also takes the values $+1$ or -1 , and which is a function of the data, adjusted by means of the statistical model in such a way that the two error probabilities are fixed: If the correct parameter is λ^b , then $P(t^b = -1|\lambda^b = +1) = \alpha$ and $P(t^b = +1|\lambda^b = -1) = 1 - \beta$. In common statistical language this means that we are testing the hypothesis $H_0 : \lambda^b = +1$, and this hypothesis is rejected if $t^b = -1$. Then α is the *level* of the test, the probability of wrong rejection, while β is the *power* of the test, the probability of rejecting the hypothesis when you should.

Note that this is still a state of the question/ answer type, albeit in a more advanced form: The question is given by the three-vector b and the two predetermined error probabilities α and $1 - \beta$. The answer is given by $\lambda^b = +1$, say, which is the conclusion we claim if we observe $t^b = +1$. So the state must involve all the quantities b , α , β and the answer ± 1 .

Say that we have done the experiment and reported the value $+1$. Then we again will use a weighing according to the error probabilities, even though these at the outset refer to different outcomes. Thus the weighted state will be

$$E = (1 - \alpha)\frac{1}{2}(I + b \cdot \sigma) + (1 - \beta)\frac{1}{2}(I - b \cdot \sigma) = \frac{1}{2}((2 - \alpha - \beta)I + (\beta - \alpha)b \cdot \sigma). \quad (22)$$

This state corresponds to what Busch [3] and Caves et al [5] call an effect $E = \frac{1}{2}(rI + ca \cdot \sigma)$, and these effects played a crucial rôle in our proof of the Born formula. In terms of the definition of an effect, we have $r = 2 - \alpha - \beta$ and $c = \beta - \alpha$.

Again it is important that both α , β and the unit vector b (with sign) can be recovered once we know the effect matrix E .

In the (r, c) -plane, the effects are limited to the triangle with corners $(0, 0)$, $(2, 0)$ and $(1, 1)$. But note that b can be replaced by $-b$ depending upon the outcome, so the triangle obtained by taking the mirror image around the line $c = 0$ is also relevant. The first triangle corresponds exactly to the limitation imposed by the hypothesis testing interpretation proposed above:

The limitations imposed by the triangle are $c \geq 0$, corresponding to $\beta \geq \alpha$, and $c \leq 1 - c$, corresponding to $\alpha \geq 0$ and $\beta \leq 1$. The bottom line $c = 0$ corresponds to $\beta = \alpha$, a case where there is no information in the reported result. The right boundary corresponds to $\alpha = 0$, and the left boundary to $\beta = 1$.

When the two error probabilities are equal, $\alpha = 1 - \beta$, we get the mixed effects. This may be interpreted as an agreement with a general property of inference problems under symmetry (see Chapter 3 in [9]): When a prior is chosen as an invariant measure of a transitive group, then Bayesian credibility statements and confidence statements (corresponding to hypothesis testing problems) are equivalent. In the present case the prior in question is just the probability $1/2$ on each of the values -1 and $+1$. And there is only one error probability to report, in the hypothesis testing case as above, in the Bayesian case the posterior probability corresponding to the result observed. With a symmetrical prior we also have $P(t^b = +1 | \lambda^b = +1) = P(\lambda^b = +1 | t^b = +1)$ and so on in this case, so that everything that is done in this subsection for this case is consistent with what was done in the previous subsection.

And of course, when $\alpha = 0$ and $\beta = 1$, both error probabilities are 0, and we get the pure states on the Bloch sphere.

A simple hypothesis problem can be inverted by exchanging hypothesis and alternative. Assume also that the reported result is opposite of what we had above, that is, -1 instead of $+1$. As on the Bloch sphere, this corresponds to an operator with the sign of c inverted, but here also with the two error probabilities exchanged. Explicitly, the operator will be

$$E_2(r, c, b) = \frac{1}{2}((2 - r)I - cb \cdot \sigma) = I - E, \quad (23)$$

that is, $E + E_2 = I$, just as $v_1^b v_1^{b\dagger} + v_2^b v_2^{b\dagger} = I$ in the Bloch sphere case.

12.4 Generalized probability.

What is left now, is to interpret the generalized probabilities assumed to be associated with the effects in the proof of the Born formulae, and to motivate the additivity of these generalized probabilities. As there, we assume that we start with a pure state a , and consider after that the generalized probability $\pi(E)$ associated with an effect $E = E(r, c, u) = E(\alpha, \beta, u)$ as introduced above. The crucial point is that these generalized probabilities must be assumed to exist, here as a result of some symmetry considerations.

In [5], and as used in the above proof of Born's formula, the basic assumptions about the generalized probabilities is that $E_1 + E_2 + \dots \leq I$ should imply $\pi(E_1 + E_2 + \dots) = \pi(E_1) + \pi(E_2) + \dots$. My intention is to indicate a proof of the above property from the general assumptions of this paper. To this end, I will need in addition to the previous Axioms a basic symmetry assumption, as expressed below. This new Axiom states in effect that, when initial and final states are transformed with the same group transformation, then the transition probability remains the same.

Recall the group K defined in Axiom 6.

Axiom 9. *When $\lambda^{a_1} = \lambda^a k$ and $\lambda^{b_1} = \lambda^b k$ with the same group element $k \in K$, then*

$$P(\lambda^{b_1} = \lambda_j k | \lambda^{a_1} = \lambda_i k) = P(\lambda^b = \lambda_j | \lambda^a = \lambda_i).$$

Theorem 8. *From the hypothesis testing interpretation and our general assumptions we have that*

$$\sum E_i \leq I \text{ implies } \pi(\sum E_i) = \sum (\pi E_i).$$

Proof: I will carry out the proof in two steps. Let

$$E = E(r, c, u) = \frac{1}{2}(r + cu \cdot \sigma), \text{ where } \|u\| = 1, 0 \leq c \leq 1, \text{ and } c \leq r \leq 2 - c.$$

(i) *For fixed u we have that $\pi(\sum r_i, \sum c_i, u) = \sum \pi(r_i, c_i, u)$.*

In the hypothesis testing interpretation

$$r = 2 - \alpha - \beta = (1 - \alpha) + (1 - \beta) = P(t = +1 | \lambda = +1) + P(t = +1 | \lambda = -1), \quad (24)$$

where we have dropped the superindex a . Similarly

$$c = \beta - \alpha = (1 - \alpha) - (1 - \beta) = P(t = +1 | \lambda = +1) - P(t = +1 | \lambda = -1) \quad (25)$$

Imagine now the following three experimental setups with the same experimental direction a :

(1) A measurement apparatus (1), whose outcome may be formulated as a hypothesis testing problem with level α_1 , power β_1 and test variable t_1 .

(2) A similar measurement apparatus (2) for the same system with characteristics α_2, β_2, t_2 .

For these two cases we find (r_i, c_i) ($i = 1, 2$) from (24) and (25).

(3) A symmetric coin is tossed to choose between the experiments (1) and (2).

The generalized probability π for this case is the probability of some state, specified by the question given by r, c and b , or by α, β and b , and the conclusion that the ideal answer is $u = b$ or $\lambda^b = +1$, i.e., the measurement gives $t^b = +1$. Now from (24) and (25) we conclude that the (r, c) -value for experiment (3) is $\frac{1}{2}(r_1 + r_2), \frac{1}{2}(c_1 + c_2)$. Thus we conclude that

$$\pi\left(\frac{1}{2}(r_1 + r_2), \frac{1}{2}(c_1 + c_2), u\right) = \frac{1}{2}\pi(r_1, c_1, u) + \frac{1}{2}\pi(r_2, c_2, u).$$

Furthermore, by considering suitable changes in the level and power of the test, again using (24) and (25), we see that $\pi(\frac{1}{2}r, \frac{1}{2}c, u) = \frac{1}{2}\pi(r, c, u)$ if both sets of values $(\frac{1}{2}r, \frac{1}{2}c)$ and (r, c) are within the triangle defining effects. Therefore we conclude that $c_1 + c_2 \geq 0$ and $c_1 + c_2 \leq r_1 + r_2 \leq 2 - c_1 - c_2$ implies

$$\pi(r_1 + r_2, c_1 + c_2, u) = \pi(r_1, c_1, u) + \pi(r_2, c_2, u).$$

The extension to several basic experiments is straightforward.

(ii) *The general case.*

Use the same setup as in (i), but assume now that the experiments (1) and (2) are measurements in different directions b_1 and b_2 , that different test levels and test powers may be used in the decision process, but that we are using physical measurement apparatus with identical display, showing the result $t = +1$ or $t = -1$. Assume that the rotation from b_1 to b_2 is defined in a specific way, so that the two directions are in correspondence relative to the measurement display. Again let experiment (3) consist of tossing a symmetric coin to choose either experiment (1) or experiment (2).

Let λ be the ‘real’ spin component in each of the three experiments, that is in the direction of the chosen measurement display. Let t_i ($i = 1, 2, 3$) be the observed spin component in experiment i . Then from the situation we have

$$P(t_3 = +1 | \lambda = +1) = \frac{1}{2}P(t_1 = +1 | \lambda = +1) + \frac{1}{2}P(t_2 = +1 | \lambda = +1), \quad (26)$$

with a similar identity for $\lambda = -1$, and, according to (24), this implies

$$r_3 = \frac{1}{2}(r_1 + r_2), \quad (27)$$

By the symmetry Axiom 8, the probability distributions of the outcomes of experiments (1) and (2) will be the same if we imagine that they have the same final vector b_1 , but that the initial vector of the experiment (2) was rotated in the opposite direction. Seen in this way, all three experiments have the same final vector, but different initial conditions. This implies that we can use an argument similar to the one used in part (i) also here, if we take the different initial conditions into account. In effect, the conclusion from this is that we use the argument that the generalized probability for experiment (3) must be the mean of the generalized probabilities for the experiments (1) and (2). Having agreed upon this, we return to the original setup with different final vectors and with formulae for effects depending upon these final vectors.

From this argument we also conclude that it is natural to associate an effect also to the artificial experiment (3). Note that this experiment can be arranged in such a way that the person reading the measurement display does not know that the experiment is artificial, i.e., that a coin has been tossed to choose between experiments (1) and (2). Since the constants r, c and u only enter the effect E in the combinations r and cu , we only have to define new values for these quantities for the experiment (3).

Consider the expected observed spin component in experiment (1). If the real parameter λ is $+1$, this expectation will be

$$\begin{aligned} & P(t_1 = 1|\lambda = 1)b_1 - (1 - P(t_1 = 1|\lambda = 1))b_1 \\ &= -b_1 + 2P(t_1 = 1|\lambda = 1)b_1 = -u_1 + 2P(t_1 = 1|\lambda = 1)u_1. \end{aligned} \quad (28)$$

The similar expression for $\lambda = -1$ will be $u_1 - 2P(t_1 = +1|\lambda = -1)u_1$. The sum of these two expectations will be

$$2P(t_1 = 1|\lambda = +1)u_1 - 2P(t_1 = 1|\lambda = -1)u_1 = 2c_1u_1$$

by (25), which gives an interpretation of the vector c_1u_1 occurring in the effect E_1 . There is of course a similar expression for experiment (2), and since the expectation of the outcome of experiment (3) must be the mean of the outcomes of experiments (1) and (2), this experiments must be assigned an effect such that

$$c_3u_3 = \frac{1}{2}(c_1u_1 + c_2u_2). \quad (29)$$

Thus it follows from the argument in the previous paragraph:

$$\pi\left(\frac{1}{2}(r_1 + r_2), \frac{1}{2}(c_1u_1 + c_2u_2)\right) = \frac{1}{2}(\pi(r_1, c_1u_1) + \pi(r_2, c_2u_2)). \quad (30)$$

Then we can adjust error probabilities in each of the experiments to get rid of the factor $1/2$ in the same way as in the previous case. This shows that in general

$$\pi(E_1 + E_2) = \pi(E_1) + \pi(E_2) \quad (31)$$

whenever the lefthand side is defined.

Some specific cases are easily verified. Begin with the case $E + E_2 = I$, as in (23), and let $\pi(E) = \pi(r, c, b) = \pi(\alpha, \beta, b)$ have some value. Then we must have that $\pi(E_2) = \pi(2 - r, -c, b) = 1 - \pi(r, c, b)$. In particular, for mixed states $\pi(1, -c, b) = 1 - \pi(1, c, b)$, and as a particular case again, the totally uninformative state has $\pi(\frac{1}{2}I) = \frac{1}{2}$. Furthermore, $\pi(0) = 0$ and $\pi(I) = 1$.

To find explicit formulae for the generalized probability $\pi(E)$, we first look at the fact, proved by Caves et al. [5] from the additivity property now motivated by Theorem 8, that $\pi(E)$ is linear in E , that is, for fixed b , linear in r and c . Fixing $\pi(b) = \pi(1, 1, b)$ for the corresponding pure state, and taking into account the values found in the previous paragraph, this implies that

$$\pi(r, c, b) = \frac{1}{2}r + (\pi(b) - \frac{1}{2})c.$$

In terms of the level and the power of the test, this is

$$\pi(\alpha, \beta, b) = 1 - \alpha\pi(b) - \beta(1 - \pi(b)).$$

As shown in the proof of Born's formula in the previous Section, the generalized probability $\pi(b)$ takes the form $\pi(b) = \frac{1}{2}(1 + a \cdot b)$, where a is the initial pure state. This gives finally

$$\pi(r, c, b) = \frac{1}{2}(r + ca \cdot b), \quad (32)$$

or

$$\pi(\alpha, \beta, b) = 1 - \frac{1}{2}(\alpha + \beta) + \frac{1}{2}(\beta - \alpha)a \cdot b. \quad (33)$$

12.5 Probability from symmetry between largescale experiments.

Four drugs A , B , C and D are being compared with respect to the expected recovery time μ they induce on patients with a certain disease. There are relatively few patients available, so one concentrates on getting information on the sign of the difference between each μ and the mean of the others, for instance

$$\lambda^A = \text{sign}(\mu_A - \frac{1}{3}(\mu_B + \mu_C + \mu_D)).$$

We will not go into detail with the experimental design here, but assume that there is an efficient design, say, of an incomplete block type, where accurate information can be obtained about one or a few such λ 's.

Just for this illustration assume that we from some experiment have obtained very accurate information that $\lambda^A = +1$, and this is the only information we have. Then we want to perform a new experiment with level α and power β in order to test an hypothesis that λ^B also is $+1$. Can we get any prior information about the result of this from the first experimental result? Informally, since μ_A is subtracted in the expression for λ^B , we should expect a probability less than $1/2$ that $\lambda^B = +1$.

In this ideal case we can get accurate information from the theory above. There is at the outset complete symmetry between the four binary parameters λ^A , λ^B , λ^C and λ^D . Permutational symmetry can always be imagined as imbedded in some rotational symmetry. Here we can imagine rotation in 3-space, since we can consider a regular tetrahedron in this space. The perpendiculars from the corners A , B , C and D of that tetrahedron to the opposite side can then be taken to represent the parameters λ^A , λ^B , λ^C and λ^D , respectively. From what is known about regular tetrahedrons, the angle between two perpendiculars is approximately 109° , with a cosine equal to $-1/3$.

At least tentatively, this can be taken as a special case of the theory above. Thus from (33), the prior probability of obtaining the result $+1$ in the last experiment is

$$1 - \frac{1}{2}(\alpha + \beta) - \frac{1}{6}(\beta - \alpha) = 1 - \frac{1}{3}\alpha - \frac{2}{3}\beta. \quad (34)$$

For the special case of an ideal experiment with $\alpha = 0$, $\beta = 1$, the probability is just $1/3$.

It seems likely that this probability assessment also can be calculated in other ways. However, to us the very important point is that the calculation here used the following two steps:

- 1) Using symmetry arguments, the probability can be assumed to exist.
- 2) Then the formula for the probability follows from the above proof of Born's formula.

In the case of the above example, we had also the crucial fact that, even though the parameter space had more dimensions, the symmetry could be reduced to a symmetry in 3-space. On *this* space, the probability space on this space and the resulting L^2 space as a Hilbert space, the above arguments for the Born formula applies. And this gives a definite formula for the probability.

In my view, the probabilities in quantum mechanics can be argued for in a similar way using a complex Hilbert space. Crucial points in the argument are first the construction of a meaningful Hilbert space which is a proper subspace of the L^2 space over the full (total) parameter space, then the result (completed in Lemma 2 above) that the vectors of that Hilbert space stand in one-to-one correspondence with the events related to the parameters of different experiments that we are interested in.

This is a parallel to the argument in the above example. And my feeling is that a similar argument can be used in several other decision problems involving inference from one macroscopic experiment to other experiments. But this is outside the scope of the present paper.

12.6 Further issues concerning decision problems.

In the Example of the previous subsection, parameters taking only two values were considered. What is new in this example from a decision theory point of view, is that we include several potential questions about the same system in the consideration.

More complicated decision problems may perhaps at least to some extent be tackled by looking at the dichotomy $\lambda^b = \lambda_k^b$ versus $\lambda^b \neq \lambda_k^b$. Or for those who know some hypothesis testing theory: What has been treated in this section, has been just the simple Neyman-Pearson case, but this is the basis for the whole statistical theory of hypothesis testing.

In any case, it is satisfying that a discussion of decision problems related to statistical experiments can be coupled to basic issues related to a physical theory like quantum mechanics. This may point at a new unification of scientific methods.

13 Basic formulae of quantum mechanics.

Having proved Born's formula, much of the results of standard quantum mechanics follow. The results of this Section are not claimed to be new of course; the point is that they can be derived from the above formulation.

Our state concept may be summarized as follows: To the state $\lambda^a(\cdot) = \lambda_k^a$ there corresponds the state vector v_k^a , and these vectors determine the transition probabilities for perfect experiments as in (17).

Theorem 9. (i) $E(\lambda^b | \lambda^a = \lambda_k^a) = v_k^{a\dagger} T^b v_k^a$, where $T^b = \sum \lambda_j^b v_j^b v_j^{b\dagger}$.
(ii) $E(f(\lambda^b) | \lambda^a = \lambda_k^a) = v_k^{a\dagger} f(T^b) v_k^a$, where $f(T^b) = \sum f(\lambda_j^b) v_j^b v_j^{b\dagger}$.

Proof: Straightforward application of Theorem 7.

Thus, in ordinary quantum mechanical terms, the expectation of every observable in any state is given by the familiar formula.

It follows from Theorem 9(i) and from the preceding discussion that the first three rules of Isham [12], as cited in the Introduction above, taken there as a basis for quantum mechanics, are satisfied.

Now turn to non-perfect experiments. In ordinary statistics, a measurement is a probability measure $Q^\theta(dy)$ depending upon a parameter θ . Assume now that such a measurement depends upon the parameter λ^b , while the current state is given by $\lambda^a = \lambda_k^a$. Then as in Theorem 9(ii)

Theorem 10. (i) Corresponding to the experiment $b \in \mathbf{A}$ one can define an operatorvalued measure M by $M(dy) = \sum_j Q^{\lambda_j^b}(dy) v_j^b v_j^{b\dagger}$. Then, given the initial state $\lambda^a = \lambda_k^a$, the probability distribution of the result of experiment b is given by $P[dy | \lambda^a = \lambda_k^a] = v_k^{a\dagger} M(dy) v_k^a$.

(ii) These operators satisfy $M[S] = I$ for the whole sample space S , and furthermore $\sum M(A_i) = M(A)$ for any finite or countable sequence of disjoint elements $\{A_1, A_2, \dots\}$ with $A = \cup_i A_i$.

Theorem 10(ii) is easily checked directly.

A more general state assumption is a Bayesian one corresponding to this setting. From Theorem 10(ii) we easily find:

Theorem 11. Let the current state be given by probabilities $\pi(\lambda_k^a)$ for different values of λ_k^a . Then, defining $\rho = \sum \pi(\lambda_k^a) v_k^a v_k^{a\dagger}$, we get $P[dy] = \text{tr}[\rho M(dy)]$.

These results are the basis for much of quantum theory, in particular for the quantum statistical inference in Barndorff-Nielsen et al. [1]; for a formulation, see also Isham [12].

Note that the density matrix $v_k^a v_k^{a\dagger}$ is equivalent to the pure state v_k^a ; similarly, a density matrix $v_j^b v_j^{b\dagger}$ is equivalent to the statement that an ideal measurement giving $\lambda^b = \lambda_j^b$ just has been performed. By straightforward application of Born's formula one gets

Theorem 12. Assume an initial state v_k^a , and assume that an ideal measurement of λ^b has been performed without knowing that value. Then this state is described by a density matrix $\sum_j |v_k^{a\dagger} v_j^b|^2 v_j^b v_j^{b\dagger}$.

This is the celebrated and much discussed projection postulate of von Neumann. Writing $P_j = v_j^b v_j^{b\dagger}$ and $\rho = v_k^a v_k^{a\dagger}$ here, the j 'th term in the last formula can be written $P_j \rho P_j$, which is a special case of the Dirac-von Neumann formula [15].

Thus the so-called von Neumann measurements can be viewed as ordinary statistical measurements in our modelling approach.

In general we have assumed for simplicity in this Section that the state vectors are nondegenerate eigenvectors of the corresponding operators, meaning that the parameter λ^a contains all relevant information about the system. This can be generalized, however.

14 Conclusion.

Although this paper contains fairly much substance, it could also be regarded as a starting point of a wider and more complete theory. Already now it goes pretty far towards giving a foundation for treating continuous variables like positions and momenta. Also, a theory for composite systems follows easily. A very natural task would be to complete the extension of the theory to non-perfect experiments in order to treat real measurement apparatus. Several other extensions can be imagined.

An important future task will be to try to discuss entanglement under this umbrella. It seems clear that, even if it should be possible to define some sort of total parameter for this case, no posterior probability could ever be defined on this total parameter. Such probabilities should be confined to results of experiments, and, as pointed out by Hess and Philipp [11] and others, even the existence of joint probabilities connected to pairs of experiments do not imply that probabilities on the full parameter space exist.

A general first goal for a theory like this could be to satisfy the requirements by Volovich [15], but we can also imagine developments beyond that. But I repeat my most important conclusion: The basis sketched here can be interpreted in a non-formal way, so it can be understood intuitively; furthermore, it can be related to ordinary statistical thinking, and thus has a link to methodology used in other empirical sciences.

There are several other approaches to quantum mechanics in the literature, but the only other really non-formal approach that I know about, is the one by Hardy [8], who defines a state as a list of probability distributions associated with every outcome of every possible experiment. There should of course be relationships between these different ways of viewing the world, but to discuss that, is beyond the scope of the present paper.

Acknowledgement.

I am grateful to Richard Gill for numerous discussions on the topics of this paper, discussions that have given me a strong feeling that I am not the only person

in this world who is interested in trying to bridge the gap between scientific traditions in this way.

References

- [1] Barndorff-Nielsen, O.E., Gill, R.D. and Jupp, P.E.: On quantum statistical inference. *J. Royal Statist. Soc. B* 65, 775-816, 2003.
- [2] Barut, A.O. and Raczka, R.: *Theory of Group Representations and Applications*. Polish Scientific Publishers, Waszaw 1977.
- [3] Busch, P.: Quantum states and generalized observables: A simple proof of Gleason's theorem. *Phys. Rev. Letters* 91 (12) 2003.
- [4] Cartwright, N.: *How the Laws of Physics Lie*. Clarendon Press, Oxford 1983.
- [5] Caves, C.M., Fuchs, C.A., Manne, K. and Renes, J.M.: Gleason-type derivations of the quantum probability rule for generalized measurements. *arXiv:quant-ph/0306179* 2003.
- [6] Freund, J.E.: *Mathematical Statistics with Applications*. Pearson Education, Upper Saddle River, NJ 2004.
- [7] Gleason, A. Measures on closed subspaces of a Hilbert space. *Journal of Mathematics and Mechanics* 6, 885-893 1957.
- [8] Hardy, L.: Why is nature described by quantum theory? In: Barrow, J.D., Davies, C.W. and Harper, C.L.: *Science and Ultimate Reality. Quantum Theory, Cosmology and Complexity*. Cambridge University Press 2004.
- [9] Helland, I.S.: *Steps Towards a Unified Basis for Scientific Models and Methods*. Book under preparation.
- [10] Helland, I.S.: Extended statistical modelling under symmetry: The link towards quantum mechanics. To appear in *Ann. Statistics* *arXiv:quant-ph/0503214* 2005.
- [11] Hess, K. and Philipp, W.: Bell's theorem: Critique and proofs without inequalities. *arXiv:quant-ph/040015* 2004.
- [12] Isham, C.J.: *Lectures on Quantum Theory*. Imperial College Press, London 1995.
- [13] Lehmann, E.L. and Casella, G.: *Theory of Point Estimation*. Springer, New York 1998.
- [14] Searle, S.R.: *Linear Models*. Wiley, New York 1971.

- [15] Volovich, I.V.: Seven principles of quantum mechanics. arXiv:quant-ph/0212126 2002.
- [16] Wolbarst, A.B.: Symmetry and Quantum Systems. Van Nostrand, New York 1977.